

Scaling of level statistics at the disorder induced metal-insulator transition*

I. Kh. Zharekeshev[†] and B. Kramer

1. Institut für Theoretische Physik, Universität Hamburg, Jungiusstrasse 9, D-20355 Hamburg, Germany

(February 1, 2008)

Abstract

The distribution of energy level separations for lattices of sizes up to $28 \times 28 \times 28$ sites is numerically calculated for the Anderson model. The results show one-parameter scaling. The size-independent universality of the critical level spacing distribution allows to detect with high precision the critical disorder $W_c = 16.35$. The scaling properties yield the critical exponent, $\nu = 1.45 \pm 0.08$, and the disorder dependence of the correlation length.

PACS numbers: 71.30.+h, 05.45.+b, 72.15.Rn

Typeset using REVTeX

*to appear in Phys. Rev. B **51**, 17239 (1995)

The statistical properties of the energy spectra of disordered quantum systems in the metallic regime can be described by the random-matrix theory initiated by Wigner and Dyson [1–3]. This was shown by analytical approaches such as the supersymmetric nonlinear σ -model formalism [4] and the impurity diagram technique [5]. A characteristic property in this regime is the repulsion between adjacent energy levels which is caused by a pronounced overlap of delocalized one-electron states.

By increasing the fluctuations of the random potential in three-dimensional (3D) systems a delocalization-localization transition is induced, the Anderson metal-insulator transition (MIT) [6]. In the insulating phase, when the magnitude of the fluctuations of the potential is much larger than the critical value, the states of the system are localized. The corresponding energy levels are not correlated due to the vanishing of their spatial overlap. Therefore, spectral fluctuations are governed by the statistics of independent random variables. A central question in discussion of the level statistics in disordered systems is how the spectral fluctuations change their behavior when extended states are transformed into localized ones when the disorder is increased.

According to the one-parameter scaling theory [7] the conductance of the system shows critical behavior at the MIT. Since the conductance is related to the spectral two-level correlation function one can expect that the level statistics also experiences the crossover between the metallic and the insulating regimes [8]. Correlations of the energy levels are contained in the distribution of the distances between successive energy levels, $P(s)$. In the metallic region $P(s)$ can be well described by the Wigner surmise

$$P_W(s) = \frac{\pi}{2} s \exp\left(-\frac{\pi}{4}s^2\right), \quad (1)$$

proposed primarily for the Gaussian orthogonal ensemble of random matrices whose elements fluctuate normally around zero with the equal variances [1,3]. Here the distances between neighboring energy levels s are measured in units of the mean level spacing Δ . In the localized regime the spacings are distributed according to a Poisson law

$$P_P(s) = \exp(-s). \quad (2)$$

The crossover from the metallic to the insulating regime is accompanied by a change in the level spacing distribution from correlated to completely uncorrelated behavior. One can, therefore, suspect that the properties of $P(s)$ for finite systems, when investigated as a function of the size of the system, would allow for the detection of the MIT and even the critical properties of the transition. As a consequence, the crossover of $P(s)$ between the Wigner (1) and the Poisson (2) statistics was extensively studied both analytically [9,10] and numerically [8,11–14]. By using of the finite-size scaling argument for $P(s)$ a new criterion for the location of the metal-insulator transition was proposed by Shklovskii *et al* [11]. This argument was cross-checked by numerically determining level statistics, and investigating the scaling properties for large s . One advantage of this method is that one needs only the energy spectrum without calculating eigenfunctions or the conductivity.

In this paper we focus again on the critical behavior of the level statistics near the disorder-induced MIT. We study the finite-size scaling properties $P(s)$ in the whole range of spacings. For the calculation of the energy levels, numerical diagonalization of the Anderson model was performed by using a modified Lanczos algorithm for systems of sizes up to $28 \times 28 \times 28$ lattice sites. We determine very precisely the critical value of the disorder W_c and the correlation length exponent ν . By considering the integrated distribution of spacings in various regions of $P(s)$, we find $W_c = 16.35$ and $\nu = 1.45 \pm 0.08$ in good agreement with similar earlier calculations [14] and previous findings based on the transfer-matrix (TM) method for the modulus of the Green's function [15,16]. We provide an independent determination of the correlation length as a function of the disorder, $\xi(W)$. Our results demonstrate unambiguously and strikingly, that the integrated level spacing distribution plays not only the role of a scaling variable at the Anderson transition, but can also be used to define qualitatively and accurately the critical properties.

The Anderson model for a disordered system is described by the Hamiltonian $H = \sum_n \varepsilon_n c_n^\dagger c_n + \sum_{n \neq m} (c_n^\dagger c_m + c_n c_m^\dagger)$, where $c_n^\dagger$ and c_n are the creation and annihilation operators of an electron at a site n in a lattice; m denotes the number Z of the nearest neighbors of the site n ($Z = 6$ in a 3D simple cubic lattice). The site energies ε_n measured in units

of the overlap integral between adjacent sites correspond to the random potential. They are uniformly distributed in the interval $[-W/2, W/2]$, where W specifies the degree of the disorder of the system. It was found earlier by the TM-method [15] that the critical disorder of the MIT which occurs in the middle of the band $E = 0$ corresponds to $W_c \approx 16.5$. Thus, for a finite system when W decreases from W_c , the level statistics should approach (1), and when $W \gg W_c$ it should obey (2) [8]. In order to find the electron spectrum in the critical region for W close to W_c we diagonalized numerically the real symmetric matrices of the Anderson Hamiltonian for simple cubic lattices of the size L^3 with periodic boundary conditions. One has to consider system sizes much larger than the lattice constant. This is necessary in order to get rid of the microscopic peculiarities of the discrete lattice model, and to be able to compare numerical with analytical results obtained in the thermodynamic limit. Despite H is represented by a very sparse matrix the computation of the energy levels of large systems is highly nontrivial due to the huge matrix size which grows as L^6 .

We have developed a considerably improved Lanczos algorithm for the eigenvalue problem of very sparse and big matrices. One of the improvements is based on the hierarchical structure of H which consists of three levels of complexity depending on the dimensionality of the lattice. Another feature is that all non-zero off-diagonal matrix elements are equal to unity. This enabled us to decrease substantially the number of multiplications during the recursion [17]. In addition, the absence of degeneracy of the eigenvalues leads to a faster convergence of the eigenvectors. We exploited the effect of level repulsion in the range of $s \lesssim 0.1$ for diminishing the number of steps of the Sturm sequence in the bisection method that is used for searching the eigenvalues of the tridiagonal matrix produced after the recursion. In this case one can govern the precision conditions for resolving two very close ‘real’ eigenvalues provided that the absolute accuracy for recognizing ‘spurious’ eigenvalues is kept as high as of the order of 10^{-12} . All of these optimizations enabled us to create a computer program which consumes much less time than a conventional implementation of the Lanczos method [17]. As a result, we succeeded in diagonalizing matrices of sizes up to 21952×21952 . We applied the algorithm to cubes of linear sizes $L = 6, 8, 12, 16, 20, 24, 28$

at various degrees of the disorder close to W_c .

One consequence of the fact that the distribution of ε_n is box-like is that the correlation length ξ depends only very weakly on the energy near the band center [16]. This follows from the phase diagram of localization which shows that $W_c(E)$ is almost constant when the mobility edge varies from $E = -6$ to 6. This allows us to use a wide central part of the band containing half of all of the eigenvalues of H , without mixing localized and delocalized states. One can check that the level statistics $P(s)$ for narrower energy intervals do not change considerably. The calculations were carried out for ensembles of different random configurations. For a given pair of $\{L, W\}$ the number of samples was chosen such that a total of approximately 10^5 eigenvalues were taken into account. Since the energy interval is so wide, the density of states $\rho = (\Delta L^3)^{-1}$ depends on energy, though only slightly, one has to unfold the spectrum. The unfolding was performed by using parabolic spline fits for $\rho(E)$.

As an example, Fig. 1 displays the histograms of $P(s)$ for $L = 6$ and 28 at various disorder W near the MIT. By increasing W the spacing distributions for both L vary continuously from $P_W(s)$ to $P_P(s)$, particularly in the range of $s \gtrsim 2$. Results for other, intermediate, sizes confirm this smooth crossover of $P(s)$. However, the behavior of the crossover depends substantially on the size of the cube. One can observe in Fig. 1 that $P(s)$ changes from (1) to (2) faster for $L = 28$ than for $L = 6$. As the size of the system increases the function $P(s)$ approaches the above two limiting regimes within a narrower interval of W around W_c . The size-dependence of $P(s)$ is observed on both sides of the MIT. Close to $W = 16.5$, the spacing distribution has almost the same form for all considered L , from 6 up to 28. The independence of $P(s)$ on L at $W = W_c$ is consistent with the prediction of a novel universal level statistics at $L \rightarrow \infty$ that exists exactly at the transition [11]. One can expect that in the thermodynamic limit there are three asymptotic shapes, namely, the Wigner surmise (1) for $W < W_c$; the Poisson law (2) for $W > W_c$, and the critical distribution $P_c(s)$ when $W = W_c$.

Since $P(s)$ scales differently with L for various regions of s , it is more convenient to

consider the integrated probability $I(s) = \int_s^\infty P(s') ds'$, which equals a fraction of those spacings which are larger than s . Because all spacings are positive, $I(0) = 1$. In addition, $\int_0^\infty I(s) ds = 1$ due to normalization to the total number of spacings in a given energy interval. Eqs. (1) and (2) give $I_W(s) = \exp(-\pi s^2/4)$ and $I_P(s) = \exp(-s)$ for the two limiting regimes. For numerical purposes we introduce $J(s) = (I(s) - I_W(s))/(I_P(s) - I_W(s))$. It describes the relative deviation of $I(s)$ from $I_W(s)$. In order to extract the dependence only on W and L and to minimize numerical errors we define a certain value of the spacing s_o for $J(s)$. The choice of the average spacing, $s_o = 1$, is not favorable, because $J(\Delta)$ exhibits a very small size effect. One can see in Fig. 1 that $P_W(s)$ and $P_P(s)$ cross at $s_1 = 0.473$ and $s_2 = 2.002$. Therefore we studied $\alpha(W, L) \equiv J(s_1)$ representative for the small- s part of $P(s)$, and $\gamma(W, L) \equiv J(s_2)$ representative for its tail, where $I_W(s)$ and $I_P(s)$ are significantly different. The former reflects the “strength” of the repulsion of two consecutive levels when the separation between them is less than s_1 , while the latter is the exponentially small probability to find a gap in the spectrum which is wider than two mean spacings Δ . For $L \rightarrow \infty$ both, α and γ , are equal to zero for $W < W_c$ and equal to unity for $W > W_c$.

Fig. 2 shows the disorder dependences of α and γ . The curves for different sizes cross at one point, W_c , which corresponds to the disorder-induced MIT. Near the critical point, as long as $|W - W_c| \lesssim 2$, $\alpha(W)$ and $\gamma(W)$ depend stronger on W for larger cubes. A similar behavior of γ was observed earlier [11] for much smaller cubes than ours. A different variable, $\int_{s_2}^\infty I(s) ds + s_2 - 1$, was studied in [14], which corresponds to using $P(s)$ in the range of large spacings, as γ does. A comparison of the two plots in Fig. 2 shows that near the MIT the scaling variables α and γ behave almost identically. The critical disorder and the scaling properties of the integrated level spacing distribution do not depend on the choice of s_o . We have checked this s_o -independence by repeating the calculations for $s_o=0.1$ and 4. However, the results for s_1 yields the highest accuracy. Using the data for α one can determine very accurately the critical value of the disorder, $W_c = 16.35 \pm 0.15$.

In the following we discuss the critical behavior of α . It characterizes the slope of the linearly growing part of $P(s)$. Assuming the validity of the one-parameter scaling $\alpha(W, L) =$

$f(L/\xi(W))$, one can replot the data of Fig. 2 by rescaling L . We numerically chose values of ξ by shifting the logarithm of the length separately for each W . An overlap between data for adjacent values of W allows to fit most of the points onto a common curve with two branches, one growing ($W > W_c$) and one decaying ($W < W_c$) (Fig. 3). Thus we find a function $\xi(W)$ up to an arbitrary factor ξ_0 . This arbitrariness can be avoided. The points for the maximum disorder $W = 20$ in Fig. 3 except of the sizes $L = 6$ and 8 correspond to the localized regime $\xi \lesssim L$. Supposing that the statistics of the levels of localized states is not sensitive to the dimensionality of system, one can match ξ for the above disorder to the relevant values of the localization length calculated by the TM-method in the quasi-1D case [15]. The function $\xi(W)$, as found by this procedure, is plotted in the inset of Fig. 3. Our values of ξ coincide with those found by the TM-method [15,16]. The agreement is slightly better for the insulating than for the metallic side.

The next step is to find the critical exponent. Using the singularity of the correlation length near W_c

$$\xi(W) = l \left| \frac{W - W_c}{W_c} \right|^{-\nu}, \quad (3)$$

where the scale l is of order of the lattice constant, one can expand the scaling function $f(L/\xi)$ into a power series $\alpha(W, L) = \alpha_c(L) + A(W - W_c)L^{1/\nu}$. By applying a conventional χ^2 -criterion for the verification of the linearized scaling hypothesis with several parameters to fit the data in Fig. 2 we found $\nu = 1.45 \pm 0.08$. It is well consistent with the critical exponent $\nu \simeq 1.53$ obtained recently by the TM-method [18]. A slightly smaller value of $\nu \simeq 1.34$ for the large- s region of $P(s)$ has been computed in [14] by using the level-statistics method similar to ours. The dependence $\xi(W)$ from (3) with $\nu = 1.45$ is shown in Fig. 3 for comparison. We note that the critical exponent could be also estimated by fitting the data for $\xi^{-1}(W)$ to the power-law function without linearizing of $\alpha(W)$ near W_c . However the relative numerical errors become markedly larger similar as in the earlier approach [15]. The accuracy of ν can then be improved by considering a narrower interval of W near the critical point or by using larger systems.

In conclusion, we have studied the integrated level spacing distribution as a scaling variable in order to detect the critical behavior of the statistics of the one-electron levels at the disorder-induced metal-insulator transition. The Anderson model was used to calculate the spectra of disordered cubic lattices of sizes varying from 6^3 up to 28^3 sites. Diagonalization of the Hamiltonian was performed numerically by means of the Lanczos algorithm especially optimized for sparse symmetric matrices with a hierarchic structure. We calculated the critical distribution $P_c(s)$ at the transition and confirmed its size-independence conjectured in [11]. Near the critical point the finite-size scaling properties of the probability $I(s)$ of neighboring spacings were examined. For the small- s and the large- s parts of $P(s)$ this probability exhibits the transition between the Wigner and the Poisson distributions which corresponds to the delocalization-localization transition. Using small spacings the critical disorder was found, $W_c = 16.35$, for the ‘box distribution’ of the site energies. Finally, we determined the disorder dependence of the correlation length $\xi(W)$ which is in good agreement with previous results obtained by the transfer-matrix method [16]. The present calculation confirms one-parameter scaling by using a completely independent method, and that the critical exponent of the purely disorder-induced metal-insulator transition is $\nu = 1.45 \pm 0.08$.

We thank B. I. Shklovskii, T. Brandes, M. Schreiber and A. Mac Kinnon for discussions. Financial support from DAAD during the stay of I. Kh. Zh. at the University of Hamburg is gratefully acknowledged. This work was supported by EU via grants SCC-CT90-0020 and CHRX-CT93-0126.

REFERENCES

- [†] Permanent address: Kazakh State University, Almaty 480121, Kazakhstan, CIS.
- [1] E. P. Wigner, Ann. Math. **62**, 548 (1955); **65**, 203 (1957).
 - [2] F. J. Dyson, J. Math. Phys. **3**, 140 (1962); **3**, 1199 (1962).
 - [3] M. L. Mehta, *Random matrices* (Academic Press, Boston, 1991).
 - [4] K. B. Efetov, Adv. Phys. **32**, 53 (1983).
 - [5] B. L. Altshuler and B. I. Shklovskii, Zh. Eksp. Teor. Fiz. **91**, 220 (1986), [Sov. Phys. JETP **64**, 127 (1986)].
 - [6] P. A. Lee and T. V. Ramakrishnan, Rev. Mod. Phys. **57**, 287 (1985).
 - [7] E. Abrahams, P. W. Anderson, D. C. Licciardello, and T. V. Ramakrishnan, Phys. Rev. Lett. **42**, 673 (1979).
 - [8] B. L. Altshuler, I. Kh. Zharekeshev, S. A. Kotochigova, and B. I. Shklovskii, Zh. Eksp. Teor. Fiz. **94**, 343 (1988) [Sov. Phys. JETP **67**, 625 (1988)]; I. Kh. Zharekeshev, Fiz. Tverd. Tela **31**, 118 (1989) [Sov. Phys. Solid State **31**, 65 (1989)].
 - [9] U. Sevan and Y. Imry, Phys. Rev. B **35**, 6074 (1987).
 - [10] A. G. Aronov, V. E. Kravtsov, and I. V. Lerner, Pis'ma v ZhETF **59**, 39 (1994), [JETP Lett. **59**, 40 (1994)].
 - [11] B. I. Shklovskii, B. Shapiro, B. R. Sears, P. Lambrianides, and H. B. Shore, Phys. Rev. B **47**, 11487 (1993).
 - [12] S. N. Evangelou and E. N. Economou, Phys. Rev. Lett. **68**, 136 (1992); S. N. Evangelou, Phys. Rev. B **49**, 16805 (1994).
 - [13] E. Hofstetter and M. Schreiber, Phys. Rev. B **48**, 16979 (1993).

- [14] E. Hofstetter and M. Schreiber, Phys. Rev. B **49**, 14726 (1994); I. Varga, E. Hofstetter, M. Schreiber, and J. Pipek, Phys. Rev. B **53**, (1994), to be published.
- [15] A. Mac Kinnon and B. Kramer, Z. Phys. B **53**, 1 (1983).
- [16] B. Kramer and A. Mac Kinnon, Rep. Prog. Phys. **56**, 1469 (1994).
- [17] J. K. Cullum and R. K. Willoughby, *Lanczos Algorithms for Large Symmetric Eigenvalue Problems* (Birkhauser, Boston, 1985).
- [18] A. Mac Kinnon, J. Phys.: Condens. Matter **6**, 2511 (1994).

FIGURES

FIG. 1. Level spacing distributions $P(s)$ for cubes of sizes $L^3 = 6^3$ and 28^3 for various values of the disorder W near the transition. Continuous curves are the Wigner surmise, Eq. (1), and the Poisson distribution, Eq. (2).

FIG. 2. Scaling variables $\alpha(W)$ and $\gamma(W)$ near W_c for different $L = 6(\diamond)$, $8(+)$, $12(\square)$, $16(\times)$, $20(\triangle)$, $24(\star)$, $28(\circ)$.

FIG. 3. Scaling variable α as a function of $L/\xi(W)$, showing one-parameter scaling. $W = 12(\diamond)$, $14(+)$, $15(\square)$, $16(\times)$, $16.5(\triangle)$, $17(\star)$, $18(\circ)$, and $20(\bullet)$. Inset: correlation length ξ as function of the disorder W . Present method ($\bullet$), TM-method ($\diamond$) [15]; $(+)$ [16]. Continuous curve: plot of Eq. (3) with $W_c = 16.35$, $\nu = 1.45$, and $l = 1$.





